



REFERENCES

PAIN AND THE BRAIN

- Aaron LA, Buchwald D. Chronic diffuse musculoskeletal pain, fibromyalgia and co-morbid unexplained clinical conditions. *Best Pract Res Clin Rheumatol* 2003;17:563–574
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision*. Washington DC: American Psychiatric Association; 2000
- Anderberg UM, Marteinsdottir I, Theorell T, et al. The impact of life events in female patients with fibromyalgia and in female healthy controls. *Eur Psychiatry* 2000;15:295–301
- Arnold LM. Biology and therapy of fibromyalgia: new therapies in fibromyalgia. *Arthritis Res Ther* 2006;8:212
- Arnold LM, Goldenberg DL, Stanford SB, et al. Gabapentin in the treatment of fibromyalgia: a randomized, double-blind, placebo-controlled, multicenter trial. *Arthritis Rheum* 2007;56:1336–1344
- Arnold LM, Hudson JI, Hess EV, et al. Family study of fibromyalgia. *Arthritis Rheum* 2004;50:944–952
- Arnold LM, Hudson JI, Keck PE Jr. Reply to causes of familial aggregation of fibromyalgia: comment on the article by Arnold et al. *Arthritis Rheum* 2004;50:3060
- Arnold LM, Hudson JI, Keck PE Jr, et al. Comorbidity of fibromyalgia and psychiatric disorders. *J Clin Psychiatry* 2006;67:1219–1225
- Arnold LM, Keck PE Jr, Welge JA. Antidepressant treatment of fibromyalgia: a meta-analysis and review. *Psychosomatics* 2000;41:104–113
- Arnold LM, Lu Y, Crofford LJ, et al. A double-blind, multicenter trial comparing duloxetine with placebo in the treatment of fibromyalgia patients with or without major depressive disorder. *Arthritis Rheum* 2004;50:2974–2984
- Arnold LM, Rosen A, Pritchett YL, et al. A randomized, double-blind, placebo-controlled trial of duloxetine in the treatment of women with fibromyalgia with or without major depressive disorder. *Pain* 2005;119:5–15
- Arthritis Foundation. *Fibromyalgia*. Atlanta, Ga: Arthritis Foundation, Inc.; 2003. Available at: http://www.foothillsfibro.org/Handouts/Fibromyalgia_brochure.pdf. Accessed Mar 31, 2008
- Bair MJ, Robinson RL, Eckert GJ, et al. Impact of pain on depression treatment response in primary care. *Psychosom Med* 2004;66:17–22
- Bair MJ, Robinson RL, Katon W, et al. Depression and pain comorbidity. *Arch Intern Med* 2003;163:2433–2445
- Basbaum AI, Fields HL. Endogenous pain control mechanisms: review and hypothesis. *Ann Neurol* 1978;4:451–462
- Bridges KW, Goldberg DP. Somatic presentation of DSM III psychiatric disorders in primary care. *J Psychosom Res* 1985;29:563–569
- Bromberger JT, Wisner KL, Hanusa BH. Marital support and remission of treated depression: a prospective pilot study of mothers of infants and toddlers. *J Nerv Ment Dis* 1993;181:40–44
- Busch AJ, Barber KAR, Overend TJ, et al. Exercise for treating fibromyalgia syndrome. *Cochrane Database Syst Rev* 2002;4:CD003786
- Crofford LJ, Rowbotham MC, Mease PJ, et al. Pregabalin for the treatment of fibromyalgia syndrome: results of a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2005;52:1264–1273
- Gallagher RM, Verma S. Managing pain and comorbid depression: a public health challenge. *Semin Clin Neuropsychiatry* 1999;4:203–220
- Geerlings SW, Twisk JWR, Beekman ATF, et al. Longitudinal relationship between pain and depression in older adults: sex, age and physical disability. *Soc Psychiatry Psychiatr Epidemiol* 2002;37:23–30
- Gendreau RM, Thorn MD, Gendreau JF, et al. Efficacy of milnacipran in patients with fibromyalgia. *J Rheumatol* 2005;32:1975–1985
- Goldenberg DL, Simms RW, Geiger A, et al. High frequency of fibromyalgia in patients with chronic fatigue seen in a primary care practice. *Arthritis Rheum* 1990;33:381–387
- Goldstein DJ, Lu Y, Detke MJ, et al. Duloxetine vs. placebo in patients with painful diabetic neuropathy. *Pain* 2005;116:109–118
- Greco T, Eckert G, Kroenke K. The outcome of physical symptoms with treatment of depression. *J Gen Intern Med* 2004;19:813–818



REFERENCES

PAIN AND THE BRAIN

- Judd LL, Akiskal HS, Paulus MP. The role and clinical significance of subsyndromal depressive symptoms (SSD) in unipolar major depressive disorder. *J Affect Disord* 1997;45:5–18
- Keller MB. Past, present, and future directions for defining optimal treatment outcomes in depression: remission and beyond. *JAMA* 2003;289:3152–3160
- Kirmayer LJ, Robbins JM, Dworkind M, et al. Somatization and the recognition of depression and anxiety in primary care. *Am J Psychiatry* 1993;150:734–741
- Kroenke K, Jackson JL, Chamberlin J. Depressive and anxiety disorders in patients presenting with physical complaints: clinical predictors and outcome. *Am J Med* 1997;103:339–347
- Kroenke K, Mangelsdorff D. Common symptoms in ambulatory care: incidence, evaluation, therapy, and outcome. *Am J Med* 1989;86:262–266
- Nicassio PM, Radojevic V, Weisman MH, et al. A comparison of behavioral and educational interventions for fibromyalgia. *J Rheumatol* 1997;24:2000–2007
- O'Malley PG, Jackson JL, Santoro J, et al. Antidepressant therapy for unexplained symptoms and symptoms syndromes. *J Fam Pract* 1999;48:980–990
- Ohayon MM. Specific characteristics of the pain/depression association in the general population. *J Clin Psychiatry* 2004;65(suppl 12):5–9
- Papakostas GI, Thase ME, Fava M, et al. Are antidepressant drugs that combine serotonergic and noradrenergic mechanisms of action more effective than the selective serotonin reuptake inhibitors in treating major depressive disorder? a meta-analysis of studies of newer agents. *Biol Psychiatry* 2007;62:1217–1227
- Park DC, Glass JM, Minear M, et al. Cognitive function in fibromyalgia. *Arthritis Rheum* 2001;44:2125–2133
- Richelson E. The clinical relevance of antidepressant interaction with neurotransmitter transporters and receptors. *Psychopharmacol Bull* 2002;36:133–150
- Rowbotham MC, Goli V, Kunz NR, et al. Venlafaxine extended release in the treatment of painful diabetic neuropathy: a double-blind, placebo-controlled study. *Pain* 2004;110:697–706
- Stahl SM. Does depression hurt? *J Clin Psychiatry* 2002;63:273–274
- Vlaeyen JW, Teeken-Gruben NJ, Goossens ME, et al. Cognitive-educational treatment of fibromyalgia: a randomized clinical trial, pt I: clinical effects. *J Rheumatol* 1996;23:1237–1245
- Weissman MM, Pilowsky DJ, Wickramaratne PJ, et al. Remissions in maternal depression and child psychopathology: a STAR*D-child report. *JAMA* 2006;295:1389–1398
- White KP, Speechley M, Harth M, et al. Comparing self-reported function and work disability in 100 community cases of fibromyalgia syndrome versus controls in London, Ontario. *Arthritis Rheum* 1999;42:76–83
- Williams DA, Cary MA, Groner KH, et al. Improving physical functional status in patients with fibromyalgia: a brief cognitive behavioral intervention. *J Rheumatol* 2002;29:1280–1286
- Wolfe F, Ross K, Anderson J, et al. The prevalence and characteristics of fibromyalgia in the general population. *Arthritis Rheum* 1995;38:19–28
- Wolfe F, Smythe HA, Yunus MB, et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia: report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990;33:160–172